

Computer Science Department

TECHNICAL REPORT

On a Conjecture of Micha Perles

Nagabhushana Prabhu

Technical Report 533

November 1990

NEW YORK UNIVERSITY



Department of Computer Science
Courant Institute of Mathematical Sciences
251 MERCER STREET, NEW YORK, N.Y. 10012

NYU COMPSCI TR-533
Prabhu, Nagabhushana
On a conjecture of Micha
Perles.



On a Conjecture of Micha Perles

Nagabhushana Prabhu

Technical Report 533

November 1990

On a Conjecture of Micha Perles

Nagabhushana Prabhu¹

Courant Institute of Mathematical Sciences

251 Mercer Street

New York, NY 10012

November 26, 1990

¹Supported in part by NSA grant MDA 904-89-H-2030, ONR grant N00014-85-K-0046 and by NSF grants CCR-8901484, CCR-8902221, CCR-8906949 and CCR-8702271

Abstract

We prove a conjecture of Micha Perles concerning simple polytopes, for a subclass that properly contains the duals of stacked and crosspolytopes. As a consequence of a special property of this subclass it also follows that, the entire combinatorial structure of a polytope in the subclass can be recovered from its graph, by applying our results recursively.

1 Introduction

Let P be a simple d -polytope and $G(P)$ the graph (1-skeleton) of P . Perles conjectured that every $(d - 1)$ -regular, induced, connected and non-separating subgraph of $G(P)$ determines a facet of P [2]. In this paper we prove the conjecture for a proper subclass of simple polytopes.

The motivation for our results comes from two subclasses of simplicial polytopes, namely the *stacked polytopes* and the *crosspolytopes*. Polytopes obtained from a simplex by successive addition of pyramids over facets are called *stacked polytopes*. Stacked polytopes form an important subclass of simplicial polytopes in that, only they attain equality in the lower bound theorem [3]. Secondly, if $\{\epsilon_1, \dots, \epsilon_d\}$ is a set of linearly independent vectors in R^d then $X = \text{conv}\{\pm\epsilon_1, \dots, \pm\epsilon_d\}$ is called a *d-crosspolytope*. d -crosspolytopes can be formed by successively building bipyramids $d - 1$ times starting with a 1-simplex.

Consider the dual analogues of the operations of stacking and forming bipyramids. It can be shown [3] that, if P is a simplicial polytope and P^* a simple polytope dual to P , then a polytope obtained by forming a pyramid over a facet of P is dual to a polytope obtained by truncating the corresponding vertex of P^* . Also, [4] any bipyramid with basis P is dual to any prism with basis P^* .

Against this background, we show that if P is any polytope for which Perles' conjecture is true, then the conjecture is also true for a polytope obtained by truncating a vertex of P and also for any prism with basis P . Since Perles' conjecture is

trivially true for a simplex, one concludes that it is true for any polytope obtained from a simplex by building prisms and truncating vertices finitely many times, in any arbitrary order. The class \mathcal{C} of polytopes so generated, properly contains duals of crosspolytopes and duals of stacked polytopes. We also show that \mathcal{C} is a proper subclass of simple polytopes. Moreover the class \mathcal{C} has the interesting property that every face of a polytope in \mathcal{C} also belongs to \mathcal{C} . This allows one to recover the entire combinatorial structure of a polytope in \mathcal{C} from its graph, by applying our results recursively.

2 Notation

Please refer to [3] for a discussion on polytopes and for related terminology. Here we merely indicate the convention that we will follow in the sequel.

If P is a polytope then its vertex set will be denoted $V(P)$ and its graph $G(P)$. The vertex set and the edge set of a graph Γ will be denoted $V(\Gamma)$ and $E(\Gamma)$ respectively. Given a d -polytope P , let $\tilde{P} = P + \mathbf{z}$ be a translate of P where \mathbf{z} is a non-zero vector in R^{d+1} . Then the convex hull of P and \tilde{P} is called a *prism with basis P* denoted $Pr(P)$. A hyperplane H in R^d is said to truncate a vertex x of P , if x and $V(P) \setminus \{x\}$ lie in different open half-spaces of H . We denote by $Tr(P)$ the intersection of P with the closed half-space containing $V(P) \setminus \{x\}$. For our purposes, it does not matter which vertex of P is truncated to obtain $Tr(P)$. $Tr(P)$ represents a (not necessarily unique) polytope obtained by truncating some

vertex of P .

Define a subclass \mathcal{C} of simple polytopes as follows: *A polytope P belongs to \mathcal{C} iff there is a sequence of polytopes*

$$P_0, P_1, \dots, P_n = P$$

where P_0 is a k -simplex ($k \geq 1$) and for $1 \leq i \leq n$ either $P_i = Tr(P_{i-1})$ or $P_i = Pr(P_{i-1})$.

From the definition of the subclass \mathcal{C} it follows that it contains the duals of stacked and crosspolytopes.

3 Perles' Conjecture for the Subclass \mathcal{C}

Theorem 1 *If Perles' conjecture is true for a simple d -polytope P then it is also true for any prism $Pr(P)$ with basis P .*

Proof : Let $Pr(P)$ be the convex hull of P and its translate $\tilde{P} = I + \mathbf{z}$. Then every vertex $v \in V(P)$ is adjacent to the vertex $\tilde{v} = v + \mathbf{z}$ of \tilde{P} . If $X \subseteq V(P)$ then \tilde{X} will denote the corresponding subset of $V(\tilde{P})$. If Λ is any induced subgraph of P then $\tilde{\Lambda}$ will denote the subgraph of $G(\tilde{P})$ induced by the corresponding vertices of \tilde{P} .

Let Γ be a d -regular, induced, connected and non-separating subgraph of $Pr(P)$. We show that Γ must determine a facet of $Pr(P)$.

If Γ is a subgraph of $G(P)$ (resp. $G(\tilde{P})$), since both Γ and $G(P)$ (resp. $G(\tilde{P})$) are d -regular graphs, $\Gamma = G(P)$ (resp. $\Gamma = G(\tilde{P})$). Hence Γ determines a facet of

$Pr(P)$. So we may assume that $V(\Gamma) \cap V(P) \neq \emptyset$ and $V(\Gamma) \cap V(\tilde{P}) \neq \emptyset$.

Let Γ_P and $\Gamma_{\tilde{P}}$ be the restrictions of Γ to P and \tilde{P} respectively. Consider any vertex v of Γ_P . v is adjacent to only one vertex in \tilde{P} . Also, Γ is d -regular. Hence v has at least $d - 1$ neighbors in Γ_P . We consider two cases.

Case 1 : Each vertex in Γ_P has exactly $d - 1$ neighbors in Γ_P .

Observe that by symmetry each vertex of $\Gamma_{\tilde{P}}$ is also $(d - 1)$ -valent in $\Gamma_{\tilde{P}}$ and that the two subgraphs Γ_P and $\Gamma_{\tilde{P}}$ are copies of each other. We also observe that:

1. Γ_P is $(d - 1)$ -regular.
2. If Γ_P has more than one connected component, pick one and call it C . Then the subgraph Γ_C of Γ induced by $V(C) \cup V(\tilde{C})$ is d -regular and hence not connected to $\Gamma \setminus \Gamma_C$ contrary to the assumption that Γ is connected. Hence Γ_P must be connected.
3. Suppose $x, y \in V(P)$ are separated by Γ_P . Let $C(x)$ and $C(y)$ be the connected components of $G(P) \setminus \Gamma_P$ containing x and y respectively. It is easy to see that $\tilde{C}(x)$ and $\tilde{C}(y)$ are separated by $\Gamma_{\tilde{P}}$ in $G(\tilde{P})$. Then Γ would separate $C(x)$ and $C(y)$, contrary to our assumption. Hence Γ_P cannot separate $G(P)$.

Since Perles' conjecture is true for P , using 1, 2 and 3 we conclude that Γ_P determines a facet F of P . Since $\Gamma_{\tilde{P}}$ is the image of Γ_P it also determines the facet \tilde{F} of \tilde{P} . So Γ determines a facet of $Pr(P)$.

Case 2 : At least one vertex in Γ_P has d neighbors in Γ_P .

Let X be the set of all the d -valent vertices in Γ_P , i.e.,

$$X = \{w \mid w \in V(\Gamma_P) \text{ and } w \text{ has } d \text{ neighbors in } \Gamma_P\}$$

Let Y be the set of all vertices in Γ_P that are adjacent to at least one vertex in X , i.e.,

$$Y = \{w \mid w \notin X; \exists x \in X, (w, x) \in E(\Gamma_P)\}$$

Since vertices in Y are $(d-1)$ -valent in Γ_P , $\tilde{Y} \subset V(\Gamma_{\tilde{P}})$. In $G(\tilde{P})$, all edges coming out of \tilde{X} terminate in \tilde{Y} . In other words, any edge path in $G(Pr(P))$ between a vertex $x \in \tilde{X}$ and a vertex $v \notin \tilde{X}$ must contain a vertex in \tilde{Y} . We know that there is a vertex $v \in (G(P) \setminus \Gamma)$ because $G(P)$ being d -regular cannot be a proper subgraph of Γ which is also d -regular (recall that we assumed $\Gamma \cap V(\tilde{P}) \neq \emptyset$.) So $\tilde{Y} \subset V(\Gamma)$ separates v and \tilde{X} contrary to the assumption that Γ does not separate $G(Pr(P))$. Hence Case 2 is impossible.

The above argument, shows that $Pr(P)$ satisfies Perles' conjecture if P does. \diamond

Theorem 2 *If Perles' conjecture is true for a simple d -polytope P , then it is also true for the d -polytope $Tr(P)$ obtained by truncating a vertex of P .*

Proof : Assume that vertex $v \in V(P)$ was truncated to obtain $Tr(P)$. Suppose $(v, w_1), \dots, (v, w_d)$ are the d edges incident on v in P . Then z_1, \dots, z_d are the new vertices in $Tr(P)$ where z_i is the intersection of (v, w_i) and the truncating hyperplane H . Also, the new facet of $Tr(P)$ (namely $H \cap Tr(P)$) is a $(d-1)$ -simplex determined by the vertex set $Z = \{z_1, \dots, z_d\}$.

Let Γ be a $(d - 1)$ -regular, connected, induced and non-separating subgraph of $G(Tr(P))$.

If $Z \cap V(\Gamma) = \emptyset$, there is nothing to prove. Also, if $Z \subseteq V(\Gamma)$ then since Z induces a $(d - 1)$ -regular subgraph $Z = V(\Gamma)$ and hence Γ determines a facet of $Tr(P)$. So the only case left to consider is where Γ contains a proper nonempty subset of Z . Since Γ is $(d - 1)$ -regular, if it contains a vertex of Z , it must contain at least $d - 2$ of its neighbors in Z . Therefore at most one vertex of Z can be left out and without loss of generality we assume $z_d \notin V(\Gamma)$. Hence, $w_i \in V(\Gamma)$ for $1 \leq i \leq d - 1$.

Consider the subgraph Γ' of $G(P)$ induced by the vertex set $(V(\Gamma) \setminus Z) \cup \{v\}$. Γ' is a $(d - 1)$ -regular, connected, induced subgraph. z_d has only one neighbour in $V(Tr(P)) \setminus V(\Gamma)$. Therefore $V(\Gamma) \cup \{z_d\}$ does not separate $G(Tr(P))$ which means Γ' does not separate $G(P)$. So Γ' determines a facet F of P and w_1, \dots, w_{d-1} are the neighbors of v in F . Let H be the supporting hyperplane for F in P . $H \cap Tr(P)$ is a facet of $Tr(P)$ and the graph of this facet is Γ ; that completes the proof. \diamond

As an immediate consequence of theorems 1 and 2 we have,

Corollary 1 *Perles' conjecture is true for every polytope in the subclass \mathcal{C} .*

The subclass \mathcal{C} has the property that any face of a polytope in \mathcal{C} also belongs to \mathcal{C} .

We prove this property in the following lemma.

Lemma 1 *If $Q \in \mathcal{C}$ and F is a facet of Q then $F \in \mathcal{C}$.*

Proof : Let $Q \in \mathcal{C}$ be a polytope for which the lemma is true. Let X be a facet of $Pr(Q)$. If $X = Q$ or $X = \tilde{Q}$ then by assumption $X \in \mathcal{C}$. If however $X = Pr(F)$ where F is a facet of Q , then since the lemma is true for Q , $F \in \mathcal{C}$ and hence $Pr(F) = X \in \mathcal{C}$. So, if the lemma is true for a $Q \in \mathcal{C}$ then it is also true for $Pr(Q)$.

Now we consider $Tr(Q)$. Let $v \in V(Q)$ be truncated to obtain $Tr(Q)$ and let H be the truncating hyperplane. Assume that H^+ contains $Tr(Q)$ (H^+ denotes one of the closed half-spaces of H). Let Y be a facet of $Tr(Q)$.

If $Y = H \cap Tr(Q)$ then Y is a simplex and hence $Y \in \mathcal{C}$. So assume $Y = F \cap H^+$ where F is a facet of Q . Since the lemma is true for Q , $F \in \mathcal{C}$. Hence $Tr(F) = Y \in \mathcal{C}$. The only case left to consider is when $Y = F$ where F is a facet of Q . Once again $Y \in \mathcal{C}$.

Therefore if the lemma is true for a $Q \in \mathcal{C}$ it is also true for $Tr(Q)$; that completes the proof.◊

As an immediate consequence of this lemma we obtain

Corollary 2 *The entire combinatorial structure of a polytope $P \in \mathcal{C}$ can be determined from $G(P)$ by repeated application of theorems 1 and 2 and lemma 1.*

The following lemma shows that \mathcal{C} is properly contained in the class of simple polytopes.

Lemma 2 *\mathcal{C} is a proper subclass of simple polytopes.*

Proof : We show that \mathcal{C} does not contain a simple 4-polytope with 9 vertices.

Suppose it did. Then, to construct the polytope we can either start with a simple 3-polytope or a 4-simplex. Suppose we started with a 4-simplex which has 5 vertices. In this case we may only truncate vertices. But each truncation (when $d=4$) increases the vertex count by 3; so we get 4-polytopes with 5, 8, 11, \dots vertices but not with 9 vertices. On the contrary suppose we started with a 3-polytope. Since constructing a prism doubles the vertex count we can only construct a prism over a 3-polytope with 4 vertices. The same argument as before shows that again we cannot obtain a 4-polytope with 9 vertices.

Now consider $C(6,4)$ - the cyclic 4-polytope with 6 vertices has 9 facets. (refer to [3] for details). It is a simplicial polytope. Its dual which is simple has 9 vertices and is hence not in \mathcal{C} . \diamond

Also, it is easy to show that the dual-stacked and the dual-crosspolytopes form proper subclasses of \mathcal{C} .

4 Remarks

Perles' conjecture is true for any simple 3-polytope[5]. So we could as well start with any simple 3-polytope and build prisms and truncate vertices finitely many times. The foregoing results would still be valid without any modification for a polytope so obtained.

References

- [1] R. Blind and P. Mani. *On puzzles and polytope isomorphisms*; Aequationes Mathematicae **34**, 287-297, 1987.
- [2] G. Kalai. *A simple way to tell a simple polytope from its graph*; Jour. Comb. Theory, Series A 49, 381-383, 1988.
- [3] A. Brøndsted. *An introduction to convex polytopes*; Springer-Verlag, 1983.
- [4] P. McMullen and G.C. Shephard. *Convex polytopes and the upper bound conjecture*; London Math. Soc. Lecture Notes, vol. 3, London: Cambridge Univ. Press, 1971.
- [5] V. Klee and P. Kleinschmidt. *Polytopal complexes and their relatives*; IMA preprint series # 384, Jan. 1988.

5.1

10

[illegible][illegible]

